



THE FACTORS CAUSING ISCHEMIC HEART DISEASE AMONG THE POPULATION OF THE SOUTHERN ARAL SEA REGION HEVE BEEN STUDIED

Ismailov K.Y.

Urgench branch of Tashkent Medical Academy

<https://orcid.org/0009-0002-0588-9438>

Abstract. The incidence and mortality of cardiovascular diseases (CVD) in Uzbekistan continue to grow and remain high. Only multifactorial preventive measures, early identification of risk factors (RF) and individuals predisposed to CVD, and changes in the lifestyle of the population through active and reasoned propaganda activities can change the situation in a positive direction. One of the factors contributing to the development of cardiovascular diseases is excess body weight and obesity. It is known that obesity is one of the most common chronic diseases in the world and leads to the development of cardiovascular diseases. Early detection and elimination of risk factors for cardiovascular disease can reduce morbidity and mortality. This article examines the development of coronary heart disease in overweight and obese adult patients by interpreting the results of electrocardiography and body weight by measuring weight (kg) and height (cm²).

Keywords: ischemic heart disease, arterial hypertension, heart failure, comorbidity, risk factors, overweight, obesity.

Introduction. According to the World Health Organization, more than one billion people are overweight, of which 300 million suffer from obesity. This increase may reach 1 billion people by 2030 (2). According to the results of a study conducted jointly with the World Health Organization (WHO) among the population of the Republic of Uzbekistan aged 18-64 years, half of the population is overweight, and 20% are obese (3). Overweight and obesity contribute to the development of cardiovascular disease (CVD) in general and coronary heart disease (CHD) in particular in part by their association with traditional and nontraditional CVD risk factors. Obesity is also considered to be an independent risk factor for CVD (1,4,6,9). An increase in body weight of 1 kg increases the risk of developing cardiovascular diseases by 3,1% . The risk of developing arterial hypertension (AH) in obese patients is 3 times higher than in people with normal body weight. There is abundant



epidemiologic evidence of an association between both overweight and obesity and CHD (1,4,5,7,8).

Purpose of the study. The purpose of the study is to investigate the role of overweight and obesity in the development of coronary heart disease among the population living in the Southern Aral Sea region of Uzbekistan.

Material and methods. 95 patients admitted to the Scientific and Practical Medical Center of Cardiology and Cardiac Surgery of the Aral Sea region were examined. According to the age classification adopted by WHO, the world is divided into age groups of 18-44, 45-59, 60-74 and 75-90 years.

The incidence of coronary heart disease in different age groups was analyzed: The patients' weight was measured in kilograms and their height in centimeters. Body mass index, developed by the Belgian sociologist and statistician Adolphe Quetelet, was calculated in kg/m^2 .

A commonly accepted body mass index (BMI) was used to determine overweight. The ratio of weight (kg) to height (cm^2) was obtained. TVI-18.5 to 24.9 kg/m is normal, 25.0 to 29.9 kg/m is overweight, and body mass index above 30.0 is considered obese. Social-hygienic and sanitary-statistical methods were used in the analysis of the collected materials.

The patients' blood pressure was measured using a tonometer. According to the latest recommendations of the European Society of Hypertension and the European Society of Cardiology, blood pressure levels are distinguished. To detect cardiovascular diseases, a 6-channel electrocardiograph with a touch screen EKG-RB600 was used. The patient underwent a standard resting electrocardiographic study and was interpreted.


Results. Of the 95 patients, 43 were men and 52 were women. 25 of them live in the city and 70 in the countryside. 4 people aged 18-44, 30 people aged 45-59, 58 people aged 60-74, and 3 people 75-90 (table).

Distribution of patients by age and gender, absolute number (%).

Age, years	Total patients	Floor		Ratio w/m
		woman	man	
18-44	4 (4,2)	2 (3,8)	2(4,6)	1:1
45-59	30 (31,57)	5 (9,7)	25 (58,2)	1:4,4
60-74	58 (61)	43 (82,7)	15 (34,9)	2,2:1
75-90	3 (3,15)	2(3,8)	1(2,3)	2:1
Total	95	52 (45,2)	43 (54,8)	1,1:1

The analysis of the occurrence of coronary heart disease by age showed that, it was observed most often in patients aged 60-74 years (61%). It was followed by 45-59-year-olds (31,57%) and the least among 18-44-year-olds (4,2%). It can be seen that the number of people suffering from ischemic heart disease increases with age. It should be noted that in the age range of 45-59 years, men made up the majority-58,2%. In contrast, women in the 60-74 age group made up the majority-82,7%.

Cases of overweight and obesity among the studied group were analyzed. The obtained results showed that all patients suffering from coronary heart disease are overweight or obese (Fig. 1). In particular, 26% have excess body weight, 38% have I-degree obesity, 11% have II-degree obesity, 11% have III-degree obesity, and in 14% IV degree of obesity was determined.



Also, 28,57% have stage 2 hypertension, and 42,86% have stage 3 hypertension. 42,86% of patients had 1 degree of arterial hypertension, 14,29% had 2 degrees, 14,29%

had 3 degrees. It is important to note that stage I hypertension was not detected in any patient.

In 57.14%, angina pectoris of the 3rd functional class was detected, in 4.76%, angina pectoris of the 2nd functional class was determined. One patient had coronary artery stenosis and coronary artery stenosis, coronary artery atherosclerosis in 4 patients, coronary artery atherosclerosis in 3 patients. Myocardial infarction with Q wave was detected in two patients. Ischemic cardiomyopathy was diagnosed in 2 patients, idiopathic pulmonary artery hypertension and tricuspid insufficiency were detected in 1 patient. During an electrocardiographic study, sinus rhythm was regular in 76.19% of cases, tachycardia - in 4.76%, extra systole - in 4.76%, atrial fibrillation - in 14.29%. When analyzing the risk level of coronary disease, 66.67% of patients were identified as risk 4, 28.57% as risk 3 and 4.76% as risk 2.

During an electrocardiographic study, sinus rhythm was regular in 76.19% of cases, tachycardia - in 4.76%, extra systole - in 4.76%, atrial fibrillation - in 14.29%. When analyzing the risk level of coronary disease, 66.67% of patients were identified as risk 4, 28.57% as risk 3 and 4.76% as risk 2 (Fig.2).




Fig. 2. Analysis of risk factors for coronary heart disease, %.

Discussion. It is clear from this that the conditions of excess body weight and obesity led to the formation of ischemic heart disease and later comorbid conditions.

In a number of studies, obesity is considered a risk factor for the development of cardiovascular diseases, and the presence of obesity in patients with ischemic heart disease contributes to its development and increased mortality. There is abundant epidemiologic evidence of an association between both overweight and obesity and



CHD (1,4,5,6,10). It can be seen that the number of people suffering from coronary heart disease increases with age.

Therefore, it was noted that the studied patients had comorbidity, and it was noted that coronary heart disease and hypertension. This indicates that patients did not pay attention to the first clinical signs due to overweight and obesity and did not consult a doctor.

It should be noted that in our studies, it was shown that overweight and obesity caused the development of ischemic heart disease and hypertension (37.5%). It can be seen that the number of people suffering from coronary heart disease increases with age. The reason for the low number of people aged 75-90 may be due to the fact that there are few people who have reached this age. Comorbid conditions were also analyzed during the study. 1.25% of the studied 95 patients suffer from the I stage of hypertension, 25.0% from the II stage, 61.25% from the III stage.

Conclusion:

1. Overweight and obesity have been found to increase the risk of developing cardiovascular diseases in the population of the South Aral Sea. There is a connection between the state of obesity and the development of cardiovascular diseases.

2. The development of ischemic heart disease among the population of the South Aral Sea can be prevented by eliminating excess body weight and obesity. This requires individual work with each person suffering from overweight and obesity by family clinic doctors.

References.

1. Вербовой А. Ф., Пашенцева А. В., Шаронова Л. А. Ожирение и сердечно-сосудистая система // Клини. мед.- 2017.- №1.-С. 95.
2. Исмаилов К. Я. Ишемическая болезнь сердца. –Ургенч, 2023.- С. 110-112.
3. Маматов А.У., Орозматов Т. Т., Мадаминов Ж. Б. Ожирение и риск развития сердечно-сосудистых заболеваний: взгляд на современную проблему // The Sci. Heritage.- 2021.-№64.- С. 36-41.
4. Санитарно-эпидемиологический комитет Республики Узбекистан. 2021г. <https://t.me/sanepidcommittee/11002>.
5. Camm A.J., Lip G.Y., De Caterina R., et al. 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation--developed with the special contribution of the European Heart Rhythm Association. // Europace. 2012. Vol. 14 №10.- P. 385-413.



6. Ismailov. K. Y. The importance of risk factors in the development of ischemic disease and death // Eurasian J.Acad.Res.- 2022. №2.-P. 189-192.

7. Ismailov, K. Y., & Djabbarova, Z. I. (2024). OBESITY AND ISCHEMIC HEART DISEASE. European Journal of Modern Medicine and Practice, 4(11), 364-367.

8. Ismailov, K. Y., & Abdullaev, I. K. Research of the Distribution and Transmission of Covid-19 Infection. JournalNX, 686-691.

9. McGill H.C., McMahan C.A., Herderick E.E., et al. Obesity accelerates the progression of coronary atherosclerosis in young men // Circulation.-2002.- Vol. 105 (23). - P. 2712–2718.

10. Katta.N., Loethen T., Lavie C. J., Alpert M.A. Obesity and Coronary Heart Disease: Epidemiology, Pathology, and Coronary Artery Imaging. <https://doi.org/10.1016/j.cpcardiol.2020.100655>.